CTP-SCU/2016011

Closed String Tachyon Driving $f(R)$ Cosmology

Peng Wang[a],* Houwen Wu[a,b],† and Haitang Yang[a]‡

[a] *Center for Theoretical Physics, College of Physical Science and Technology, Sichuan University, Chengdu, 610064, PR China and*

[b] *Center for the Fundamental Laws of Nature, Harvard University, Cambridge, MA, 02138, USA*

arXiv:1610.03952v1 [hep-th] 13 Oct 2016

Abstract

To study quantum effects on the bulk tachyon dynamics, we replace R with $f(R)$ in the low-energy effective action that couples gravity, the dilaton, and the bulk closed string tachyon of bosonic closed string theory and study properties of their classical solutions. The α' corrections of the graviton-dilaton-tachyon system are implemented in the $f(R)$. We obtain the tachyon-induced rolling solutions and show that the string metric does not need to remain fixed in some cases. The singular behavior of more classical solutions are investigated and found to be modified by quantum effects. In particular, there could exist some classical solutions, in which the tachyon field rolls down from a maximum of the tachyon potential while the dilaton expectation value is always bounded from above during the rolling process.

*Electronic address: pengw@scu.edu.cn

†Electronic address: wu˙houwen@g.harvard.edu

‡Electronic address: hyanga@scu.edu.cn

Contents

I. Introduction 2

II. Coupled System of Fields 4

III. Tachyon-Driven Rolling Solutions 6

A. Analytic at R_0 7

1. $R_0 = 0$ 8

2. $F(R_0) = 0$ 9

B. Pole at R_0 10

1. $R_0 = 0$ 10

2. $R_0 \neq 0$ 11

IV. Cosmological Solutions 12

A. Constant H 12

B. Constant T 15

V. Conclusion 18

A. Appendix 19

Acknowledgments 21

References 21

I. INTRODUCTION

In the standard cosmological model, the Friedmann equations are derived from Hilbert action, coupling the metric with ad hoc matter sources. In string theory, a low energy effective action arises from the requirement of quantum conformal invariance. The tree level of this action couples the metric, the dilaton and the axion. This action, implying the dynamics of gravitational field and other background fields, has been used in many areas. One of the most important applications is to develop a string cosmology [1, 2].

There exist two ways to improve this string effective action [3, 4]. The first one is α'-controlled expansion, which includes higher derivatives of the metric and background fields. Another one is string coupling g_s-controlled expansion (higher-genus expansion) and which reflects the higher loop string interactions. The α'-controlled expansion, which becomes significant in Planck and high curvature region, had been discussed in [5, 6]. When curvature and energy scale grow up, the usual simplest perturbative expansion of the string effective action becomes no-go theorem. In this condition, the higher order expansions should be taken into account. A good review of string cosmology is given by [7] and references therein.

The instabilities associated with open string tachyon is well understood presently. The stable vacuum is the vacuum of closed string without open string excitations and D-branes. The open string tachyon rolls down from the perturbative unstable vacuum to an excited state of closed string carrying the energy of the D-brane. On the other hand, the instabilities associated with closed string tachyon is a much more difficult problem since the action is non-polynomial. Some progresses have been achieved in the calculation of the effective potential [8–11] by truncating the polynomial to quintic order. Those calculations indicate that there do exist a local minimum and several saddle points in the closed string tachyon and dilaton effective potential. The conjecture that the action has to vanish at the closed string field theory (CSFT) vacuum, made in [8, 12], is also implied. At the closed string vacuum, parallel to the conclusions in open string theory, one is led to believe that spacetime itself ceases to exist. To make these conjectures more reliable, higher order calculations or more desirable analytical methods are necessary.

For these reasons, studying the string effective action that couples the metric, the dilaton and the tachyon is of importance. One can expect that this action will reveal some features of string theory and related problems. Some progresses have been made in recent developments. The author of [13] discussed the solutions of the effective action with tachyon and B-field and analyzed the solutions in AdS_3 background. In [14], Brandenberger *et.al.* found a nonsingular and static tachyon condensation by discussing an effective theory with a non-vanishing dilaton potential. The quantum effects of graviton-dilaton-tachyon system is investigated in [15] where it shows that the singular behavior of classical solutions should be modified by quantum effects. The author of [16] uses some constraints to fix the form of action without computing the beta functions. Some other progresses of graviton-dilaton-tachyon system refer to [17–26].

The aim of this paper is an extension of [8, 12] to study the α' corrections of the graviton-dilaton-tachyon system. The α' correction should also satisfy the quantum conformal invariance and then brings higher derivative terms of gravitational field as well as other background fields. To preserve the covariance and gauge invariance, the same expansion order of α' could lead to different effective actions [7, 27]. To simplify the story, we only implement corrections to the graviton by replacing R with $f(R)$. This could be accomplished by integrating out other massive fields in the action.

In [12], the rolling process triggered by the tachyon was investigated for tachyonic potentials. It was found that during the rolling process, the string metric did not evolve while the dilaton rolled to strong coupling. In the framework of $f(R)$ gravity, our analysis shows some interesting results. In section III, we find for tachyon-induced rolling solutions that time evolution of the string metric could be nontrivial in some cases. In section IV, assuming the Hubble parameter of the string metric is constant, we obtain a set of solutions, in which the tachyon field rolls down from the top of some tachyon potential while the string coupling is always finite during the rolling process.

This paper is organized as follows. In section II, we derive the equations of motion of graviton, dilaton, and tachyon field in $f(R)$ theories. In section III, we define tachyon-induced rolling solutions and solve the equations of motion for them. More classical solutions are discussed in section IV. The section V is our conclusion.

II. COUPLED SYSTEM OF FIELDS

The low energy effective action for the metric, the dilaton, and the tachyon is given by

$$S = \frac{1}{2\kappa^2}\int d^{d+1}x\sqrt{-g}e^{-2\Phi}\left[f(R) + 4(\nabla\Phi)^2 - (\nabla T)^2 - 2V(T)\right], \tag{1}$$

where $g_{\mu\nu}$ is string metric, Φ is dilaton, T is tachyon with potential $V(T)$, and $f(R)$ is the generalized gravity. The number of spatial dimensions is d. Since here we focus on the tachyon driving solutions, the dilaton potential is set to zero for simplicity. Varying the action (1) with respect to $g_{\mu\nu}$, Φ, and T, we find that the equations of motion for graviton,

dilaton and tachyon are

$$
\begin{aligned}
&F\left(R\right)R_{\mu\nu}-\nabla_{\mu}T\nabla_{\nu}T+4g_{\mu\nu}\left[F\left(R\right)-1\right]\left(\nabla\Phi\right)^{2}-4g_{\mu\nu}\nabla^{\alpha}\Phi\nabla_{\alpha}F\left(R\right)\\
&-2g_{\mu\nu}\left[F\left(R\right)-1\right]\nabla^{2}\Phi+g_{\mu\nu}\nabla^{2}F\left(R\right)+2F\left(R\right)\nabla_{\nu}\nabla_{\mu}\Phi\\
&-\nabla_{\nu}\nabla_{\mu}F\left(R\right)-4\left[F\left(R\right)-1\right]\nabla_{\nu}\Phi\nabla_{\mu}\Phi+2\nabla_{\mu}\Phi\nabla_{\nu}F\left(R\right)+2\nabla_{\nu}\Phi\nabla_{\mu}F\left(R\right)=0, \qquad (2)\\
&\frac{1}{2}\left[F\left(R\right)R-f\left(R\right)\right]+2\left\{d\left[F\left(R\right)-1\right]+1\right\}\left(\nabla\Phi\right)^{2}-2d\nabla^{\alpha}\Phi\nabla_{\alpha}F\left(R\right)\\
&-\left\{d\left[F\left(R\right)-1\right]+1\right\}\nabla^{2}\Phi+\frac{1}{2}d\nabla^{2}F\left(R\right)+V\left(T\right)=0, \qquad (3)\\
&\nabla^{2}T-2\nabla^{\mu}\Phi\nabla_{\mu}T-V'\left(T\right)=0, \qquad (4)
\end{aligned}
$$

where $F(R)\equiv df(R)/dR$. As in [12], we make the following ansatz

$$
\begin{aligned}
ds^{2}&=-dt^{2}+a\left(t\right)^{2}\delta_{ij}dx^{i}dx^{j},\\
\Phi&=\Phi\left(t\right), \qquad (5)\\
T&=T\left(t\right).
\end{aligned}
$$

For the string metric in eqn. (5), the Ricci scalar is

$$R=2d\dot{H}+d\left(d+1\right)H^{2}, \qquad (6)$$

where $H=\frac{\dot{a}}{a}$. In this case, the 00 and ij components of gravitational equations become

$$
\begin{aligned}
&F\left(R\right)\left(d\dot{H}+dH^{2}\right)+\dot{T}^{2}-2\ddot{\Phi}+\left\{2\left[F\left(R\right)-1\right]\dot{\Phi}-\dot{F}\left(R\right)\right\}dH=0, \qquad (7)\\
&F\left(R\right)\left(\dot{H}+dH^{2}\right)-4\left[F\left(R\right)-1\right]\dot{\Phi}^{2}+4\dot{\Phi}\dot{F}\left(R\right)+2\left[F\left(R\right)-1\right]\ddot{\Phi}\\
&+2\left[F\left(R\right)\left(d-1\right)-d\right]H\dot{\Phi}-\ddot{F}\left(R\right)-\left(d-1\right)H\dot{F}\left(R\right)=0, \qquad (8)
\end{aligned}
$$

which can be rearranged into two equivalent equations

$$
\begin{aligned}
&\frac{1}{2}F\left(R\right)\left(d-1\right)\dot{H}+\frac{1}{2}\dot{T}^{2}+2\left[F\left(R\right)-1\right]\dot{\Phi}^{2}-2\dot{\Phi}\dot{F}\left(R\right)\\
&-F\left(R\right)\ddot{\Phi}+F\left(R\right)H\dot{\Phi}+\frac{1}{2}\ddot{F}\left(R\right)-\frac{1}{2}H\dot{F}\left(R\right)=0, \qquad (9)\\
&\frac{1}{2}F\left(R\right)\left(d-1\right)dH^{2}-2d\left[F\left(R\right)-1\right]\dot{\Phi}^{2}+2d\dot{\Phi}\dot{F}\left(R\right)+\left\{d\left[F\left(R\right)-1\right]+1\right\}\ddot{\Phi}\\
&+\left\{\left[d\left(F\left(R\right)-1\right)-2F\left(R\right)+1\right]\dot{\Phi}-\frac{1}{2}\left(d-2\right)\dot{F}\left(R\right)\right\}dH-\frac{1}{2}d\ddot{F}\left(R\right)-\frac{1}{2}\dot{T}^{2}=0. \qquad (10)
\end{aligned}
$$

The equations of motion for the dilaton and the tachyon are

$$\frac{1}{2}\left[F\left(R\right)R-f\left(R\right)\right]-2\left[d\left(F\left(R\right)-1\right)+1\right]\dot{\Phi}^{2}+2d\dot{\Phi}\dot{F}\left(R\right)$$
$$+\left\{d\left[F\left(R\right)-1\right]+1\right\}\left(\ddot{\Phi}+dH\dot{\Phi}\right)-\frac{1}{2}d\left[\ddot{F}\left(R\right)+dH\dot{F}\left(R\right)\right]+V\left(T\right)=0, \tag{11}$$

$$\ddot{T}+\left(dH-2\dot{\Phi}\right)\dot{T}+V'\left(T\right)=0. \tag{12}$$

Note that there are four differential equations for three unknown dynamical variables: $a\left(t\right)$, $\Phi\left(t\right)$, and $T\left(t\right)$. Therefore, one of the four equations should be redundant. In fact, it shows in the appendix that eqns. (7), (8), and (12) could guarantee that eqn. (11) holds whenever $\dot{T}\neq 0$.

III. TACHYON-DRIVEN ROLLING SOLUTIONS

In this section, we consider a general class of potentials $V(T)$ for a tachyon T that has a local maximum at $T=0$, which can be written as

$$V\left(T\right)=V(0)-\frac{1}{2}m^{2}T^{2}+\mathcal{O}\left(T^{3}\right). \tag{13}$$

In [12] where $f\left(R\right)=R$, the rolling solutions driven by the tachyon were discussed. The ansatzes for $T\left(t\right)$ and $\Phi\left(t\right)$ were assumed to be

$$\begin{aligned} T\left(t\right) &= e^{mt}+\sum_{n\geq 2}t_{n}e^{nmt}, \\ \Phi\left(t\right) &= \sum_{n\geq 2}\phi_{n}e^{nmt}, \\ H\left(t\right) &= \sum_{n\geq 2}h_{n}e^{n\gamma t}, \end{aligned} \tag{14}$$

where $T\rightarrow 0$ for $t\rightarrow-\infty$, and $\Phi\left(t\right)$ has exponentials subleading to e^{mt} since the tachyon drives the rolling in the very early time. For the case with $f\left(R\right)=R$, eqn. (10) becomes

$$\frac{1}{2}\left(d-1\right)dH^{2}=\frac{1}{2}\dot{T}^{2}-\ddot{\Phi}+dH\dot{\Phi}, \tag{15}$$

which gives that $H\rightarrow 0$ when $t\rightarrow-\infty$, which is consistent with the ansatz for $H\left(t\right)$ proposed in eqn. (14). In fact, it showed in [12] that $H\left(t\right)$ vanished identically for the tachyon-driven rolling solutions.

For the case with a general form of $f(R)$, we build an analogous tachyon-driven rolling solution:

$$T(t) = e^{\gamma t} + \sum_{n\geq 2} t_n e^{n\gamma t},$$
$$\Phi(t) = \sum_{n\geq 2} \phi_n e^{n\gamma t}, \tag{16}$$

where γ is a positive real number, and the first term in $T(t)$ is the solution to the linearized tachyon equation of motion. The arbitrary constant multiplying the first term in $T(t)$ can be absorbed, as we did, by a redefinition of time. Unlike in the case with $f(R) = R$, eqn. (10) fails to require that $H \to 0$ when $t \to -\infty$. There might exist possible solutions with $H(t)$ which does not go to zero when $t \to -\infty$. Thus, one might consider a more general ansatz for $H(t)$:

$$H(t) = H_0 + \sum_{n\geq 1} h_n e^{n\gamma t}. \tag{17}$$

If the rolling process is solely trigger by the tachyon filed, one needs that $H_0 = h_1 = 0$. As we show below, for such ansatz nontrivial $H(t)$ could exist in some cases. However to explore more possibilities, we now do not require $H_0 = h_1 = 0$. Plugging $T(t)$ and $\Phi(t)$ from eqns. (16) into the tachyon equation (12), we find

$$\gamma^2 + dH_0\gamma - m^2 = 0, \tag{18}$$

which gives $\gamma = m\left(\sqrt{1+\frac{d^2H_0^2}{4m^2}} - \frac{dH_0}{2m}\right)$. The corresponding Ricci scalar is

$$R = R_0 + \sum_{n\geq 1} r_n e^{n\gamma t}, \tag{19}$$

where $R_0 = d(d+1)H_0^2$ and r_n depends on $h_{m\leq n}$. We assume $f(R)$ is analytic along the real axis except certain poles.

In the rest of this section, we will use eqns. (7), (8) and (12) to solve for R_0, h_n, ϕ_n and t_n. And eqn. (11) could be used to determine the value of $V(0)$. Note that in [12] where $f(R) = R$, it showed that $V(0) = 0$, $R_0 = 0$ and $h_n = 0$.

A. Analytic at R_0

Now $f(R)$ is assumed to be analytic at R_0. Thus, we can expand $f(R)$ at $R = R_0$:

$$f(R) = f(R_0) + \sum_{l\geq 1} \alpha_l (R - R_0)^l. \tag{20}$$

Eqn. (19) gives $F(R) = F(R_0) + \mathcal{O}(e^{\gamma t})$. Thus, the leading terms of the 00 and ij components of gravitational equations (7) and (8) both become $F(R_0)\, dH_0^2 = 0$. Given $R_0 = d(d+1)H_0^2$, one finds that either $F(R_0) = 0$ or $R_0 = 0$. The tachyon equation (12) is trivial at the leading order. The leading order of the dilation equation (11) gives

$$V(0) = -\frac{f(R_0)}{2}, \tag{21}$$

where we use $F(R_0)\, R_0 = 0$. Note that eqn. (21) must be satisfied to admit the tachyon rolling solution. For example, if $f(R) = R - \Lambda$ where Λ is the cosmological constant and $R_0 = 0$, one has $V(0) = \frac{\Lambda}{2}$. In what follows, we will solve for h_n, ϕ_n and t_n in the two cases with $R_0 = 0$ and $F(R_0) = 0$.

1. $R_0 = 0$

If $f(R)$ is analytic at $R = 0$, $F(R) = 1 + \sum_{l\geq 2} l\alpha_l R^{l-1}$. Plugging eqns. (16) and (17) with $H_0 = 0$ into the equations of motion, one could determine the coefficients t_n, ϕ_n and h_n order by order. Note that $\gamma = m$ for $H_0 = 0$.

At $\mathcal{O}(e^{mt})$, eqns. (7) and (8) give

$$\begin{aligned} dmh_1 e^{mt} &= 0, \\ \left(1 - 4dm^2\alpha_2\right) mh_1 e^{mt} &= 0, \end{aligned} \tag{22}$$

which leads to $h_1 = 0$. Tachyon equation is also trivial at this order. At $\mathcal{O}(e^{2mt})$,eqns. (7) and (8) give

$$\begin{aligned} \left(m^2 + 2dmh_2 - 8m^2\phi_2\right) e^{2mt} &= 0, \\ 2m\left(1 - 16dm^2\alpha_2\right) h_2 e^{2mt} &= 0, \end{aligned} \tag{23}$$

which yield

$$\begin{aligned} \phi_2 &= \frac{1}{8} + \frac{dh_2}{4m}, \\ h_2 = 0 \text{ or } \alpha_2 &= \frac{1}{16dm^2}. \end{aligned} \tag{24}$$

Note that we have $h_2 = 0$ if $\alpha_2 \neq \frac{1}{16dm^2}$. Now we prove by induction that $H(t)$ vanishes identically if $\alpha_2 \neq \frac{1}{4dN^2m^2}$ for $N = 2, 3\cdots$. First assume that $h_2 = h_3 = \cdots = h_N = 0$. Since $\dot{\Phi}(t) \sim e^{2mt}$, eqn. (8) gives

$$(N+1)\left[1 - 4d\alpha_2 (N+1)^2 m^2\right] mh_{N+1} e^{(N+1)mt} + \mathcal{O}\left(e^{(N+2)mt}\right) = 0, \tag{25}$$

which yields $\left[1-4d\alpha_2\left(N+1\right)^2m^2\right]h_{N+1}=0$. Therefore, if $\alpha_2\neq\frac{1}{4dN^2m^2}$ for $N=2,3\cdots$, $h_{N+1}=0$.

On the other hand, if $\alpha_2=\frac{1}{4dN^2m^2}$ for some positive integer $N>1$, h_N could be nonzero. Therefore, there might be solutions with nonzero $H(t)$ in such cases. Consider an example with

$$f\left(R\right)=R+\frac{R^2}{16dm^2}+\sum_{l\geq3}\alpha_lR^l \text{ and } V\left(T\right)=-\frac{1}{2}m^2T^2,$$

we find solutions up to $\mathcal{O}\left(e^{4mt}\right)$

$$\begin{aligned}T\left(t\right)&=e^{mt}+\frac{e^{3mt}}{16}+\mathcal{O}\left(e^{4mt}\right),\\ \Phi\left(t\right)&=\left(\frac{1}{8}+\frac{dh}{4m}\right)e^{2mt}+\mathcal{O}\left(e^{4mt}\right),\\ H\left(t\right)&=he^{2mt}+\mathcal{O}\left(e^{4mt}\right),\end{aligned}\tag{26}$$

where h is a free parameter. As far as the tachyon rolling ansatz is concerned, the parameter h could be arbitrary.

2. $F\left(R_0\right)=0$

Suppose $F\left(R\right)$ is analytic at R_0 where $F\left(R_0\right)=0$. Then, one can expand $F\left(R\right)$ at R_0:

$$F\left(R\right)=\sum_{l\geq1}\beta_l\left(R-R_0\right)^l,\tag{27}$$

where we assume $\beta_1\neq0$ for simplicity. Plugging eqns. (16) and (17) into eqns. (7), (8), and (12) gives the recurrence relations for h_n,ϕ_n,and t_n with necessary initial values. In fact, after putting the ansatzes into eqns. (7), (8), and (12), one has at $\mathcal{O}\left(e^{n\gamma t}\right)$ that

$$\begin{aligned}2n\gamma\left[\left(n\gamma+dH_0\right)\phi_n-d^2H_0^2\beta_1h_n\right]&=G_n\left(h_{i<n},\phi_{i<n},t_{i<n}\right),\\ 2n\gamma\left(n\gamma+dH_0\right)\phi_n+2n\gamma d\left[\left(n^2\gamma^2-H_0^2\right)+\left(d-1\right)H_0n\gamma\right]\beta_1h_n&=F_n\left(h_{i<n},\phi_{i<n},t_{i<n}\right),\\ \gamma\left[\left(n^2-1\right)\gamma+dH_0\left(n-1\right)\right]t_n&=H_n\left(h_{i<n},\phi_{i<n},t_{i<n}\right),\end{aligned}\tag{28}$$

where G_n, F_n and H_n are functions of only $h_{i<n}$, $\phi_{i<n}$, and $t_{i<n}$. Solving the above equations for h_n, ϕ_n, and t_n, we find that the recurrence relations for h_n, ϕ_n,and t_n for $n \geq 2$ are

$$\begin{aligned}
\phi_n &= \frac{G_n\left(h_{i<n}, \phi_{i<n}, t_{i<n}\right)}{2n\gamma\left(n\gamma + dH_0\right)} + \frac{dH_0^2}{n\gamma + dH_0} \frac{F_n\left(h_{i<n}, \phi_{i<n}, t_{i<n}\right) - G_n\left(h_{i<n}, \phi_{i<n}, t_{i<n}\right)}{2n^2\gamma^2\left[n\gamma + (d-1)H_0\right]}, \\
h_n &= \frac{F_n\left(h_{i<n}, \phi_{i<n}, t_{i<n}\right) - G_n\left(h_{i<n}, \phi_{i<n}, t_{i<n}\right)}{2dn^2\gamma^2\beta_1\left[n\gamma + (d-1)H_0\right]}, \\
t_n &= \frac{1}{\gamma} \frac{H_n\left(h_{i<n}, \phi_{i<n}, t_{i<n}\right)}{\left(n^2-1\right)\gamma + dH_0\left(n-1\right)}.
\end{aligned} \tag{29}$$

From eqns. (16), one obtains that $t_1 = 1$ and $\phi_1 = 0$. Since $\dot{H}$, $F(R) \sim \mathcal{O}(e^{\gamma t})$ and $\dot{T}^2, \Phi \sim \mathcal{O}(e^{2\gamma t})$,all terms of eqn. (9) except $\frac{1}{2}\ddot{F}(R)$ are at $\mathcal{O}(e^{2\gamma t})$. Given $\ddot{F}(R) = 2d\beta_1 h_1 \gamma^3 e^{\gamma t} + \mathcal{O}(e^{2\gamma t})$, one has $h_1 = 0$ since $\beta_1 \neq 0$ by assumption. After the initial conditions $h_1 = 0$, $\phi_1 = 0$, and $t_1 = 1$ are obtained, one could use the recurrence relations (29) to find values of h_n, ϕ_n,and t_n. For example, we have for $n = 2$ that

$$\phi_2 = \frac{1}{8}, h_2 = \frac{\gamma}{4d\beta_1\left[(d+1)H_0^2 - 2dH_0\gamma - 4\gamma^2\right]}, t_2 = 0, \tag{30}$$

where h_2 is generally not zero.

B. Pole at R_0

We now study the scenario in which $f(R)$ has a pole of order $|L|$ at R_0. Therefore, the Laurent series expansion of $f(R)$ at R_0 is

$$f(R) = \sum_{l \geq L} \alpha_l \left(R - R_0\right)^l, \tag{31}$$

where $\alpha_L \neq 0$ and $L \leq -1$ is some negative integer. We now consider two cases: $R_0 = 0$ and $R_0 \neq 0$.

1. $R_0 = 0$

Since $H_0 = 0$, the ansatz for $H(t)$ (17) becomes

$$H(t) = \sum_{n \geq N} h_n e^{nmt}, \tag{32}$$

where $h_N e^{Nmt}$ is assumed to be the nonzero leading term. If $H(t) \neq 0$, one has some integer $N \geq 1$ for which $h_N \neq 0$. Plugging eqn. (32) into $R = 2d\dot{H} + d(d+1)H^2$ gives for nonzero

integer l that

$$R^l = r_N^l e^{Nlmt} + \mathcal{O}\left(e^{(Nl+1)mt}\right), \tag{33}$$

where $r_N = 2dNmh_N$. Thus, we have

$$F(R) = \sum_{l\geq L} l\alpha_l R^{l-1} = Lr_N^{L-1}\alpha_L e^{N(L-1)mt} + \mathcal{O}\left(e^{[N(L-1)+1]mt}\right). \tag{34}$$

Given eqn. (34), the ij components of gravitational equation (8) becomes

$$-Lr_N^{l-1}\alpha_L N^2 (L-1)^2 m^2 e^{N(L-1)mt} + \mathcal{O}\left(e^{[N(L-1)+1]mt}\right) = 0,$$

where gives $h_N^{L-1}\alpha_L = 0$. Since $\alpha_L \neq 0$, one obtains $h_N = 0$ and hence a contradiction, which means $H(t) = 0$. However $f(R)$ blows up at $R = 0$, and hence the tachyon rolling ansatzes (16) and (17) do not solve the equations of motion (7), (8) and (12) in this case.

2. $R_0 \neq 0$

The ansatz for $H(t)$ is

$$H(t) = H_0 + \sum_{n\geq N} h_n e^{n\gamma t}, \tag{35}$$

where $h_N e^{N\gamma t}$ is the first nonzero term in the series. Using eqn. (31), one has

$$f(R) = \alpha_L r_N^L e^{NL\gamma t} + \mathcal{O}\left(e^{(NL+1)\gamma t}\right), \tag{36}$$

where $r_N = 2d\left[N\gamma + (d+1)H_0\right]h_N$. Therefore, eqn. (7) gives

$$\left[H_0 - N(L-1)\gamma\right] dH_0 L r_N^{L-1}\alpha_L e^{N(L-1)\gamma t} + \mathcal{O}\left(e^{(N(L-1)+1)\gamma t}\right) = 0. \tag{37}$$

If $N\gamma + (d+1)H_0 \neq 0$ and $H_0 - N(L-1)\gamma \neq 0$ for any positive integer N, eqn. (37) gives $\alpha_L = 0$, which means $H(t) = H_0$. However, $f(R)$ blows up at $R = R_0$, and hence the tachyon rolling ansatz does not solve the equations of motion. If $N\gamma + (d+1)H_0 = 0$ or $H_0 - N(L-1)\gamma = 0$ some N, the coefficient of $e^{N(L-1)\gamma t}$ in eqn. (37) is zero without making $\alpha_L = 0$. In principle, one could put the ansatz into the equations of motion and solve for h_n, ϕ_n, and t_n. However, one could not obtain the recurrence relations as in the $F(R_0) = 0$ case. One might need other means to find values of h_n, ϕ_n,and t_n.

IV. COSMOLOGICAL SOLUTIONS

In section III, we considered a special class of solutions to the equations of motion, whose ansatzes are given in eqns. (16) and (17). For the case with $f(R) = R$, the string metric remains fixed for such solutions. However for some more general form of $f(R)$, there might exist tachyon-driven rolling solutions with nonzero $H(t)$. It appears that the behavior of the classical solutions is richer in the $f(R)$ theory. In this section, we study the classical solutions beyond the ansatzes (16) and (17). We are still interested in the form of solutions in eqns. (5), in which the metric is the spatially flat FRW metric. The equations of motion then reduce to eqns. (7), (8), (11), and (12). Note that if $V(T) = V(-T)$, the solutions are invariant under the "time-reversal" transformations $t \to -t$, for which

$$H \to -H,\ R \to R,\ \Phi \to \Phi,\ T \to -T. \tag{38}$$

However, the equations of motion might become nonlinear higher order differential equations, which are difficult to solve. To investigate the properties of their solutions, we consider two simple scenarios, the one with constant H, and the other with constant T.

A. Constant H

If we assume $H(t) = H_0$ which is a constant, there are three independent equations of motion for $\Phi(t)$ and $T(t)$. Therefore, one of these equations determines the form of the tachyon potential $V(T)$, which depends on H_0. In other words, only some particular potentials $V(T)$ admit the classical solutions with $H(t) = H_0$. In this case, the ij components of gravitational equation (8) becomes

$$dF(R_0) H_0^2 - 4[F(R_0) - 1]\dot{\Phi}^2 + 2[F(R_0) - 1]\ddot{\Phi} + 2dH_0[F(R_0) - 1]\dot{\Phi} - 2H_0 F(R_0)\dot{\Phi} = 0, \tag{39}$$

where $R(t) = R_0 = d(d+1)H_0^2$. To solve this equation, we can introduce the variable $\tau = dH_0 t$. In what follows, the dot denotes t-derivative, and the prime denotes τ-derivative.

If $F(R_0) = 1$ and $H_0 = 0$, eqn. (39) becomes trivial and hence there are only two independent equations of motion for $\Phi(t)$ and $T(t)$. In this case, $\Phi(t)$ and $T(t)$ can be solved for any $V(T)$, and the properties of these solutions have been discussed in [12]. It has been found that the string coupling always became divergent at some time.

If $F(R_0) = 1$ and $H_0 \neq 0$, the solution to eqn. (39) is

$$\Phi(\tau) = \frac{\tau - \tau_0}{2}, \tag{40}$$

where the integration constant τ_0 is a constant time translation and can be set to zero for simplicity from now on. The dilaton Φ goes to infinity at $t = +\infty$. This solution evolves to a singular configuration with a strongly coupled background at infinite string time. The dilaton equation (11) gives $V(T) = \frac{1}{2}[f(R_0) - R_0]$. However, eqn. (7) becomes

$$\dot{T}^2 = -dH_0^2 < 0, \tag{41}$$

which contradicts the tachyon field T being real. Thus, there are no classical solutions with constant H in this case.

If $F(R_0) \neq 1$, solving eqn. (39) for Φ gives

$$\Phi_{\pm}(\tau) = \frac{1}{2}\ln\left|\frac{e^{\tau}}{1 \pm \gamma e^{A\tau}}\right| + C_{\Phi}, \tag{42}$$

where C_{Φ} is an integration constant, $A = 1 + \frac{F(R_0)}{d[F(R_0)-1]}$, and $\gamma = |1 - F(R_0)|$. Plugging $\Phi(t)$ into the dilaton equation (11), one could solve it for $V(T(\tau))$:

$$\frac{V(\tau)}{d^2H_0^2} = -\frac{F(R_0)R_0 - f(R_0)}{2d^2H_0^2} + sgn(F(R_0) - 1)\frac{\rho A e^{A\tau}}{\pm 1 + \gamma e^{A\tau}}, \tag{43}$$

where $\rho = \frac{F(R_0)[dF(R_0)-d+1]}{2d}$, and $sgn(x)$ is the sign function with $sgn(x) = \frac{x}{|x|}$. Plugging $\Phi(t)$ into the gravitational equation (7) gives

$$T_{\pm}'^2 = \alpha\frac{\pm 1 + \beta e^{A\tau}}{(1 \pm \gamma e^{A\tau})^2}, \tag{44}$$

where $B = d + F(R_0) - 3dF(R_0) + 2dF^2(R_0)$, $\alpha = \frac{F(R_0)(d+1)-d}{d}$, and $\beta = \frac{B}{2d|1-F(R_0)|}$. When $F(R_0) = \frac{d}{d+1}$, one has that $\alpha = A = 0$. In this case, T stays constant, and

$$\Phi(\tau) = \frac{\tau}{2} + C_{\Phi}, \tag{45}$$

which is the linear dilaton solution. When $F(R_0) \neq \frac{d}{d+1}$, we list the sign of $T_{\pm}'^2$ for the possible values of $F(R_0)$ in TABLE I for $d < 6$ and TABLE II for $d \geq 6$, where we define $F_{\pm} = \frac{-1+3d\pm\sqrt{1-6d+d^2}}{4d}$ such that $B = 0$ when $F(R_0) = F_{\pm}$. Note that $0 < F_- < F_+ < \frac{d}{d+1}$. When $T_{\pm}'^2 \geq 0$, we integrate eqn. (44) and obtain

$$T_{\pm}(\tau) = \frac{2\sqrt{\alpha}}{A}\,\mathrm{Re}\left[\sqrt{\frac{\gamma-\beta}{\gamma}}\mathrm{arctanh}\left(\sqrt{\frac{\gamma}{\gamma-\beta}}\sqrt{\pm 1 + \beta e^{A\tau}}\right) - \mathrm{arctanh}\left(\sqrt{\pm 1 + \beta e^{A\tau}}\right)\right] + C_T, \tag{46}$$

	$F(R_0) > \frac{d}{d+1}$	$F(R_0) < \frac{d}{d+1}$
$T_+'^2$	$T_+'^2 > 0$	$T_+'^2 < 0$
$T_-'^2$	$T_-'^2 \geq (<) 0$ for $e^{A\tau} \geq (<) \frac{2d\gamma}{B}$	$T_-'^2 \geq (<) 0$ for $e^{A\tau} \leq (>) \frac{2d\gamma}{B}$

TABLE I: The sign of $T_\pm'^2$ for $d < 6$.

	$F(R_0) > \frac{d}{d+1}$	$F_+ \leq F(R_0) < \frac{d}{d+1}$ or $F(R_0) \leq F_-$	$F_- < F(R_0) < F_+$
$T_+'^2$	$T_+'^2 > 0$	$T_+'^2 < 0$	$T_+'^2 \geq (<) 0$ for $e^{A\tau} \geq (<) \frac{2d\gamma}{\|B\|}$
$T_-'^2$	$T_-'^2 \geq (<) 0$ for $e^{A\tau} \geq (<) \frac{2d\gamma}{\|B\|}$	$T_-'^2 \geq (<) 0$ for $e^{A\tau} \leq (>) \frac{2d\gamma}{\|B\|}$	$T_-'^2 < 0$

TABLE II: The sign of $T_\pm'^2$ for $d \geq 6$.

where C_T is an integration constant. In principle, one could use eqns. (43) and (46) to find $V(T)$.

When $\tau \to -\infty$, $\Phi_\pm(\tau)$ always goes to $-\infty$. Defining $\tau = \tau_a$ such that $\gamma e^{A\tau_a} = 1$, one finds that $\Phi_-(\tau_a) = +\infty$. As $\tau \to +\infty$, one has for $\Phi_\pm(\tau)$ that

$$\Phi_\pm(\tau) \to \begin{cases} +\infty, & \text{for } A < 1 \\ C_\Phi, & \text{for } A = 1 \\ -\infty, & \text{for } A > 1 \end{cases}. \tag{47}$$

For $\Phi_+(\tau)$ with $A < 1$ and $\Phi_-(\tau)$, the string coupling always diverges at some time. However for $\Phi_+(\tau)$ with $A \geq 1$, the string coupling always stay finite. In FIG. 1(a), we plot $\Phi_+(\tau)$ for $F(R_0) = 2$. Note that $A \geq 1$ implies that $F(R_0) \leq 0$ or $F(R_0) > 1$. TABLEs I and II show that $T_+'^2 < 0$ when $F(R_0) \leq 0$. Thus when $A \geq 1$, the solution $T_+(\tau)$ only exists for $F(R_0) > 1$. When $F(R_0) > 1$, eqns. (43) and (46) shows that as $\tau \to -\infty$

$$T_+(\tau) \sim \sqrt{\alpha}\tau,\ \frac{dV(T)}{dT} \sim d^2 H_0^2 \frac{\rho A^2}{\sqrt{\alpha}} e^{A\tau},\ \text{and}\ \frac{d^2V(T)}{dT^2} \sim d^2 H_0^2 \frac{\rho A^4}{\alpha} e^{A\tau},$$

and as $\tau \to +\infty$

$$T_+(\tau) \sim C_T,\ \frac{dV(T)}{dT} \sim d^2 H_0^2 \frac{\rho A^2}{\sqrt{\alpha\beta\gamma}} e^{-A\tau/2} \sim 0,\ \text{and} \frac{d^2V(T)}{dT^2} \sim -d^2 H_0^2 \frac{\rho A^4}{2\alpha\beta} < 0,$$

where we use $\alpha > 0$, $\beta > 0$, and $\rho > 0$ for $F(R_0) > 1$. Therefore when $\tau = +\infty$, the tachyon field T stays at a maximum of $V(T)$. Since $T_+(\tau)$ goes to $-\infty$ as $\tau \to -\infty$, one finds for



(a) Plot of $\Phi_+(\tau)$ (b) Plot of $V(T)$

FIG. 1: Plots of $\Phi_+(\tau)$ and $V(T)$ for $F(R_0)=2$, where we have $C_\Phi=0$.

$T\to-\infty$ that the tachyon potential becomes

$$V(T)\sim-\frac{F(R_0)R_0-f(R_0)}{2}+\rho Ad^2H_0^2e^{AT/\sqrt{\alpha}}. \tag{48}$$

In FIG. 1(b), we plot $V(T)$ for $F(R_0)=2$. Recalling that $\tau=dH_0t$, we find for $H_0<0$ and $F(R_0)>1$, that the solution $T_+(t)$ in eqn. (46) describes the scenario in which the tachyon field is at a maximum of $V(T)$ at $t=-\infty$ and begins to roll down $V(T)$ afterwards. At $t=\infty$, $T_+(t)$ goes to $-\infty$, and $V(T)$ is given in eqn. (48). Moreover, this tachyon rolling down scenario is free of singularities since the Ricci scalar is constant, and the string coupling is always finite.

B. Constant T

If $T=T_0$ such that $V'(T_0)=0$, the tachyon equation (12) becomes trivial. As a result, one could solve the gravitational equations (7) and (8) for $H(t)$ and $\Phi(t)$. The dilaton equation (11) impose constraints on the integration constants. This scenario could provide some insights into the possible final state of bulk tachyon condensation. The case with $f(R)=R$ has been discussed in [15], where the classical solutions always were found to evolve from or to singular configurations. We here investigate the singular behavior of the solutions in the case with $f(R)=R+\frac{\alpha}{2}R^2$, in which the perturbation method is used to find solutions. The perturbation method gives how these solutions are altered for non-zero but small α. In doing so, we assume that the altered solutions can be Taylor expanded in α. In addition, it turns out that the forms of the solutions depend on the sign of $V(T_0)$.

We calculate the perturbative solutions to $\mathcal{O}(\alpha)$ for $V(T_0) = 0$ and discuss the singular behavior of the solutions for $V(T_0) > 0$ and $V(T_0) < 0$.

Substituting the Taylor expansions of $H(t)$ and $\Phi(t)$ in powers of α

$$\begin{aligned} H(t) &= H_0(t) + \alpha H_1(t) + \mathcal{O}\left(\alpha^2\right), \\ \Phi(t) &= \Phi_0(t) + \alpha \Phi_1(t) + \mathcal{O}\left(\alpha^2\right), \end{aligned} \tag{49}$$

into the gravitational equations (7) and (8), one finds

$$\begin{aligned} d\dot{H}_0 + dH_0^2 - 2\ddot{\Phi}_0 &= 0, \\ \dot{H}_0 + dH_0^2 - 2H_0\dot{\Phi}_0 &= 0, \\ d\dot{H}_1 + 2dH_0H_1 - 2\ddot{\Phi}_1 &= F(t), \\ \dot{H}_1 + 2dH_0H_1 - 2H_1\dot{\Phi}_0 - 2H_0\dot{\Phi}_1 &= G(t), \end{aligned} \tag{50}$$

where we define

$$\begin{aligned} F(t) &\equiv -d\left[R_0\left(\dot{H}_0 + H_0^2\right) + \left(2R_0\dot{\Phi}_0 - \dot{R}_0\right)H_0\right], \\ G(t) &\equiv -R_0\left(\dot{H}_0 + dH_0^2\right) + 4R_0\dot{\Phi}_0^2 - 4\dot{\Phi}_0\dot{R}_0 - 2R_0\ddot{\Phi}_0 - 2(d-1)R_0H_0\dot{\Phi}_0 + \ddot{R}_0 + (d-1)H_0\dot{R}_0. \end{aligned} \tag{51}$$

If $V(T_0) = 0$, solving eqns. (50) and using eqn. (11) to constrain the integration constants, one finds that the solutions to $\mathcal{O}(\alpha)$ are

$$\begin{aligned} H_\pm(t) &= \pm\frac{1}{\sqrt{d}(t-t_0)} + \frac{\alpha h^\pm}{4(t-t_0)^3} + \alpha c(t-t_0)^{-2}, \\ \Phi_\pm(t) &= \frac{\pm\sqrt{d}-1}{2}\ln|t-t_0| + \Phi_0 + \frac{\alpha\phi^\pm}{(t-t_0)^2} - \frac{\alpha c\left(d \mp \sqrt{d}\right)}{2(t-t_0)}, \end{aligned} \tag{52}$$

where c and Φ_0 and are integration constants, and we have

$$\begin{aligned} h^\pm &= \mp\frac{\left(\mp\sqrt{d}+1\right)^2\left(5 \pm 6\sqrt{d} + 5d\right)}{\sqrt{d}}, \\ \phi^\pm &= \frac{\left(\pm\sqrt{d}-1\right)^2\left(\pm 5d^{\frac{3}{2}} + 4d \pm \sqrt{d} - 2\right)}{16}. \end{aligned} \tag{53}$$

Integrating $H_\pm(t)$, we find

$$a_\pm(t) = a_0|t-t_0|^{\frac{\pm 1}{\sqrt{d}}}\exp\left[-\frac{\alpha h^\pm}{8(t-t_0)^2} - \frac{\alpha c}{t-t_0}\right], \tag{54}$$



(a) Plot of $a_+(t)$

(b) Plot of $\Phi_+(t)$

(c) Plot of $a_-(t)$

(d) Plot of $\Phi_-(t)$

FIG. 2: Plots of $a_\pm(t)$ and $\Phi_\pm(t)$, where we have $d=3$, $\alpha=1$, $t_0=0$, $\Phi_0=0$, $a_0=1$ and $c=0$.

where a_0 is a constant. The $\mathcal{O}(\alpha)$ corrections would not change the asymptotic behavior of the solutions at $t=\pm\infty$. Around $t=t_0$, the $\mathcal{O}(\alpha)$ corrections could dramatically change the asymptotic behavior of the these solutions. We plot $a_\pm(t)$ and $\Phi_\pm(t)$ in FIG. 2, where we have $d=3$, $\alpha=1$, $t_0=0$, $\Phi_0=0$, $a_0=1$ and $c=0$. For example, the solutions $a_+(t)$ and $\Phi_+(t)$ with $\alpha=0$ evolve from (to) big bang (big crunch) at $t=t_0$, while they have a very weakly string coupled background. However for $\alpha>0$, $a_+(t)$ goes to infinity at $t=t_0$, while the string coupling becomes divergent. Note that these perturbative solution are valid when $\alpha(t-t_0)^{-2}\ll 1$. Therefore, the higher order corrections are necessary to study the singular behavior of classical solutions. However, our analysis gives a sense of how quantum effects modify the singular behavior of classical solutions.

For $V(T_0) > 0$, the leading terms of the solutions are

$$H_0^{\pm}(t) = \pm\frac{C}{\sqrt{d}}\frac{1}{\sinh C(t-t_0)},$$
$$\Phi_0^{\pm}(t) = \frac{1}{2}\left(\pm\sqrt{d}-1\right)\ln\left|\sinh\frac{C}{2}(t-t_0)\right| - \frac{1}{2}\left(\pm\sqrt{d}+1\right)\ln\left|\cosh\frac{C}{2}(t-t_0)\right| + \Phi_0, \quad (55)$$

where $C = \sqrt{2V(T_0)}$, and t_0 and Φ_0 are integration constants. For $V(T_0) < 0$, the leading terms are

$$H_0^{\pm}(t) = \pm\frac{C}{\sqrt{d}}\frac{1}{\sin C(t-t_0)},$$
$$\Phi_0^{\pm}(t) = \frac{1}{2}\left(\pm\sqrt{d}-1\right)\ln\left|\sin\frac{C}{2}(t-t_0)\right| - \frac{1}{2}\left(\pm\sqrt{d}+1\right)\ln\left|\cos\frac{C}{2}(t-t_0)\right| + \Phi_0, \quad (56)$$

where $C = \sqrt{-2V(T_0)}$, and t_0 and Φ_0 are integration constants. These solutions have a singularity at $t = t_0$. Around $t = t_0$, their singular behaviors are the same as in the case with $V(T_0) = 0$. Therefore, the last two equations in eqns. (50) give that the singular behaviors of $H_1(t)$ and $\Phi_1(t)$ at $t = t_0$ are also the same as in the case with $V(T_0) = 0$.

V. CONCLUSION

In [12], the tachyon-induced rolling solutions have been considered using the low-energy effective field equations, which were derived from the effective action (1) with $f(R) = R$. To gain some insight into quantum effects on the tachyon dynamics, in this paper we investigated the behavior of the classical solutions of the low-energy effective action (1) of the graviton-dilaton-tachyon system, in which quantum corrections are included only in $f(R)$ for simplicity. After the equations of motion were obtained in section II, we solved them for the tachyon-induced rolling ansatzes (16) and (17) in section III. Finally, more classical solutions were discussed in section IV.

In [12] where $f(R) = R$, it showed that $H(t)$ vanished identically for the tachyon-induced rolling solutions. For more general forms of $f(R)$, we found in subsection III A that there were some cases in which $H(t)$ could be nonzero. Moreover, we solved the equations of motion assuming $H(t) = H_0$ which was a constant. When $F(R_0) = 1$ and $H_0 = 0$, the properties of classical solutions have been discussed in [12], and it has been found that the dilaton always rolled toward stronger coupling. However for $F(R_0) > 1$, we found that there existed some solutions in which the string coupling could always stay finite. In the case of

$f(R) = R + \frac{\alpha}{2}R^2$, we also solved the equations of motion assuming $T(t) = T_0$, whose scenario is related to the possible final state of bulk tachyon condensation. It turned out for some solutions that higher order terms in $f(R)$ could dramatically change their singular behavior. Since we only used an effective model, the $f(R)$ gravity theory, to investigate quantum effects on tachyon dynamics, one might not take our analysis too seriously. However, our analysis suggests that quantum corrections should be important to understand tachyon dynamics.

Appendix A: Appendix

On defining

$$G_1 \equiv F(R)\left(d\dot{H} + dH^2\right) + \dot{T}^2 - 2\ddot{\Phi} + \left[2\left(F(R) - 1\right)\dot{\Phi} - \dot{F}(R)\right]dH, \tag{A1}$$

$$G_2 \equiv F(R)\left(\dot{H} + dH^2\right) - 4\left[F(R) - 1\right]\dot{\Phi}^2 + 4\dot{\Phi}\dot{F}(R) + 2\left[F(R) - 1\right]\ddot{\Phi} + 2\left[F(R)(d-1) - d\right]H\dot{\Phi} - \ddot{F}(R) - (d-1)H\dot{F}(R), \tag{A2}$$

$$G_3 \equiv \frac{1}{2}\left[F(R)R - f(R)\right] - 2\left\{d\left[F(R) - 1\right] + 1\right\}\dot{\Phi}^2 + 2d\dot{\Phi}\dot{F}(R) + \left\{d\left[F(R) - 1\right] + 1\right\}\left(\ddot{\Phi} + dH\dot{\Phi}\right) - \frac{1}{2}d\left[\ddot{F}(R) + dH\dot{F}(R)\right] + V(T), \tag{A3}$$

$$G_4 \equiv \ddot{T} + \left(dH - 2\dot{\Phi}\right)\dot{T} + V'(T), \tag{A4}$$

the equations of motion (7), (8), (11), and (12) become $G_i = 0$. Supposing that $G_1 = G_2 = G_3 = 0$, we now show that this leads to $G_4 = 0$ whenever $\dot{T} \neq 0$.

Differentiating $G_1 = 0$ with respect to time gives

$$\dot{T}\ddot{T} = -\frac{\dot{F}(R)\left(d\dot{H} + dH^2\right)}{2} - \frac{F(R)\left(d\ddot{H} + 2dH\dot{H}\right)}{2} + \dddot{\Phi} - \left\{\left[F(R) - 1\right]\dot{\Phi} - \frac{\dot{F}(R)}{2}\right\}d\dot{H} - \left\{\left[F(R) - 1\right]\ddot{\Phi} + \dot{F}(R)\dot{\Phi} - \frac{\ddot{F}(R)}{2}\right\}dH, \tag{A5}$$

where we use $R = 2d\dot{H} + d(d+1)H^2$. By multiplying G_4 by $\dot{T}$, eliminating $\dot{T}\ddot{T}$ through

eqn. $(A5)$ and $\dot{T}^2$ through the equation of motion $G_1 = 0$, we find

$$\dot{T}G_4 - \left(dH - 2\dot{\Phi}\right) G_1 - \frac{1}{2}\frac{dG_1}{dt} = -\frac{\dot{F}(R)\left(d\dot{H} + dH^2\right)}{2} - \frac{F(R)\left(d\ddot{H} + 2dH\dot{H}\right)}{2} + \dddot{\Phi}$$
$$- \left\{\left[F(R) - 1\right]\dot{\Phi} - \frac{\dot{F}(R)}{2}\right\} d\dot{H} - \left\{\left[F(R) - 1\right]\ddot{\Phi} + \dot{F}(R)\dot{\Phi} - \frac{\ddot{F}(R)}{2}\right\} dH \qquad \text{(A6)}$$
$$- \left(dH - 2\dot{\Phi}\right)\left\{F(R)\left(d\dot{H} + dH^2\right) - 2\ddot{\Phi} + \left[2\left(F(R) - 1\right)\dot{\Phi} - \dot{F}(R)\right] dH\right\} + V'(T)\dot{T} = 0.$$

Differentiating $G_3 = 0$ with respect to time gives

$$\frac{\dot{F}(R)R}{2} - 4\left[d\left(F(R) - 1\right) + 1\right]\dot{\Phi}\ddot{\Phi} - 2d\dot{F}(R)\dot{\Phi}^2 + 2d\ddot{\Phi}\dot{F}(R) + 2d\dot{\Phi}\ddot{F}(R) + d\dot{F}(R)\left(\ddot{\Phi} + dH\dot{\Phi}\right)$$
$$+ \left\{d\left[F(R) - 1\right] + 1\right\}\left(\dddot{\Phi} + d\dot{H}\dot{\Phi} + dH\ddot{\Phi}\right) - \frac{1}{2}d\left[\dddot{F}(R) + d\dot{H}\dot{F}(R) + dH\ddot{F}(R)\right] = -V'(T)\dot{T}. \qquad \text{(A7)}$$

Differentiating $G_2 = 0$ with respect to time and multiplying by $d/2$ on both sides gives

$$d\left[F(R) - 1\right]\dddot{\Phi} - \frac{d}{2}\dddot{F}(R) - 4d\left[F(R) - 1\right]\dot{\Phi}\ddot{\Phi} - 2d\dot{F}(R)\dot{\Phi}^2 + 2d\ddot{\Phi}\dot{F}(R)$$
$$+2d\dot{\Phi}\ddot{F}(R) + d\dot{F}(R)\ddot{\Phi} + d^2H\dot{\Phi}\dot{F}(R) - \frac{d^2}{2}\dot{H}\dot{F}(R) - \frac{d^2}{2}H\ddot{F}(R) \qquad \text{(A8)}$$
$$= -\frac{d\dot{F}(R)\left(\dot{H} + dH^2\right)}{2} - \frac{dF(R)\left(\ddot{H} + 2dH\dot{H}\right)}{2} + dH\dot{\Phi}\dot{F}(R)$$
$$-d\left[F(R)(d-1) - d\right]\dot{H}\dot{\Phi} - d\left[F(R)(d-1) - d\right]H\ddot{\Phi} - \frac{d\dot{H}\dot{F}(R)}{2} - \frac{dH\ddot{F}(R)}{2}.$$

Subtracting eqn. $(A8)$ from eqn. $(A7)$ gives

$$\frac{dG_3}{dt} - \frac{d}{2}\frac{dG_2}{dt} = \dddot{\Phi} - 4\dot{\Phi}\ddot{\Phi} + \left[F(R) + 1\right]d\dot{H}\dot{\Phi} - \frac{dF(R)\ddot{H}}{2} \qquad \text{(A9)}$$
$$+\left[F(R) + 1\right]dH\ddot{\Phi} - \frac{dH\ddot{F}(R)}{2} + dH\dot{\Phi}\dot{F}(R) - d^2F(R)H\dot{H} + \frac{d\dot{F}(R)H^2}{2} + V'(T)\dot{T} = 0.$$

Subtracting eqn. $(A6)$ from eqn. $(A9)$ gives

$$\frac{dG_3}{dt} - \frac{d}{2}\frac{dG_2}{dt} - \dot{T}G_4 + \left(dH - 2\dot{\Phi}\right)G_1 + \frac{1}{2}\frac{dG_1}{dt}$$
$$= dH\left\{2\left(F(R) - 1\right)\ddot{\Phi} - \ddot{F}(R) + F(R)\left(\dot{H} + dH^2\right) - (d-1)H\dot{F}(R)\right. \qquad \text{(A10)}$$
$$\left. -4\left[F(R) - 1\right]\dot{\Phi}^2 + 4\dot{\Phi}\dot{F}(R) + 2\left[F(R)(d-1) - d\right]H\dot{\Phi}\right\}.$$

Comparing eqn. $(A10)$ with eqn. $(A4)$, we obtain the relation for G_1, G_2, G_3 and G_4

$$\left(dH - 2\dot{\Phi}\right)G_1 + \frac{1}{2}\frac{dG_1}{dt} - \frac{d}{2}\frac{dG_2}{dt} - dHG_2 + \frac{dG_3}{dt} - \dot{T}G_4 = 0, \qquad \text{(A11)}$$

which shows that $G_4 = 0$ if $G_1 = G_2 = G_3 = 0$ and $\dot{T} \neq 0$.

Acknowledgments

We are grateful to Jun Tao and Zheng Sun for useful discussions. This work is supported in part by NSFC (Grant No. 11005016, 11175039 and 11375121).

---

[1] M. Gasperini and G. Veneziano, "The Pre-big bang scenario in string cosmology," Phys. Rept. **373**, 1 (2003) [arXiv:hep-th/0207130].

[2] T. Battefeld and S. Watson, "String gas cosmology," Rev. Mod. Phys. **78**, 435 (2006) [arXiv:hep-th/0510022].

[3] M. B. Green, J. H. Schwarz and E. Witten, "SUPERSTRING THEORY. VOL. 1: INTRODUCTION," Cambridge, Uk: Univ. Pr. (1987) 469 P. (Cambridge Monographs On Mathematical Physics)

[4] J. Polchinski, "String theory. Vol. 1: An introduction to the bosonic string," Cambrigde University Press (1998).

[5] M. D. Pollock, "On the Quantum Cosmology of the Superstring Theory Including the Effects of Higher Derivative Terms," Nucl. Phys. B **324**, 187 (1989).

[6] M. D. Pollock, "On the derivation of the Wheeler-DeWitt equation in the heterotic superstring theory," Int. J. Mod. Phys. A **7**, 4149 (1992).

[7] M. Gasperini, "Elements of string cosmology," Cambridge, UK: Cambridge Univ. Pr. (2007) 552 p.

[8] H. Yang and B. Zwiebach, "A Closed string tachyon vacuum?," JHEP **0509**, 054 (2005) [arXiv:hep-th/0506077].

[9] N. Moeller and H. Yang, "The nonperturbative closed string tachyon vacuum to high level," JHEP **0704**, 009 (2007) [arXiv:hep-th/0609208].

[10] N. Moeller, "Closed bosonic string field theory at quintic order: Five-tachyon contact term and dilaton theorem," JHEP **0703**, 043 (2007) [arXiv:hep-th/0609209].

[11] N. Moeller, "Closed bosonic string field theory at quintic order. II: Marginal deformations and effective potential," JHEP **0709**, 118 (2007) [arXiv:0705.2102 [hep-th]].

[12] H. Yang and B. Zwiebach, "Rolling closed string tachyons and the big crunch," JHEP **0508**, 046 (2005) [arXiv:hep-th/0506076].

[13] T. Suyama, "Tachyon Condensation with B-field," JHEP **0702**, 050 (2007) [arXiv:hep-th/0610127].

[14] R. H. Brandenberger, A. R. Frey and S. Kanno, "Towards A Nonsingular Tachyonic Big Crunch," Phys. Rev. D **76**, 063502 (2007) [arXiv:0705.3265 [hep-th]].

[15] T. Suyama, "Quantum Effects on Tachyon Dynamics," JHEP **0808**, 051 (2008) [arXiv:0805.3563 [hep-th]].

[16] M. Headrick, "A Note on tachyon actions in string theory," Phys. Rev. D **79**, 046009 (2009) [arXiv:0810.2809 [hep-th]].

[17] S. S. Razamat, "On fluctuations of closed string tachyon solitons," JHEP **0711**, 007 (2007) [arXiv:0707.2310 [hep-th]].

[18] I. Swanson, "Cosmology of the closed string tachyon," Phys. Rev. D **78**, 066020 (2008) [arXiv:0804.2262 [hep-th]].

[19] I. Y. Aref'eva and A. S. Koshelev, "Cosmological Signature of Tachyon Condensation," JHEP **0809**, 068 (2008) [arXiv:0804.3570 [hep-th]].

[20] J. Alexandre, A. Kostouki and N. E. Mavromatos, "Tachyon-Dilaton-induced Inflation as an alpha-prime-resummed String Background," JHEP **0903**, 022 (2009) [arXiv:0811.4607 [hep-th]].

[21] Y. Hikida, R. R. Nayak and K. L. Panigrahi, "D-branes in a big bang/big crunch universe: Misner space," JHEP **0509**, 023 (2005) [arXiv:hep-th/0508003].

[22] H. Singh, "(A)symmetric tachyon rolling in de Sitter spacetime: An Universe devoid of planck density," Nucl. Phys. B **734**, 169 (2006) [arXiv:hep-th/0508101].

[23] T. Suyama, "Closed string tachyon condensation in supercritical strings and RG flows," JHEP **0603**, 095 (2006) [arXiv:hep-th/0510174].

[24] J. Genenberg and V. Suneeta, "The Fixed Points of RG Flow with a Tachyon," JHEP **0609**, 045 (2006) [arXiv:hep-th/0605230].

[25] O. Bergman and S. S. Razamat, "Toy models for closed string tachyon solitons," JHEP **0611**, 063 (2006) [arXiv:hep-th/0607037].

[26] C. Escamilla-Rivera, G. Garcia-Jimenez, O. Loaiza-Brito and O. Obregon, "Closed String Tachyon: Inflation and Cosmological Collapse," Class. Quant. Grav. **30**, 035005 (2013) [arXiv:1110.6223 [gr-qc]].

[27] R. R. Metsaev and A. A. Tseytlin, "Order alpha-prime (Two Loop) Equivalence of the String

Equations of Motion and the Sigma Model Weyl Invariance Conditions: Dependence on the Dilaton and the Antisymmetric Tensor," Nucl. Phys. B **293**, 385 (1987).

[28] A. A. Starobinsky, "A New Type of Isotropic Cosmological Models Without Singularity," Phys. Lett. B **91**, 99 (1980).

[29] S. Capozziello, "Curvature quintessence," Int. J. Mod. Phys. D **11**, 483 (2002) [arXiv:gr-qc/0201033].

[30] S. Capozziello, V. F. Cardone, S. Carloni and A. Troisi, "Curvature quintessence matched with observational data," Int. J. Mod. Phys. D **12**, 1969 (2003) [arXiv:astro-ph/0307018].

[31] S. Capozziello, S. Carloni and A. Troisi, "Quintessence without scalar fields," Recent Res. Dev. Astron. Astrophys. **1**, 625 (2003) [arXiv:astro-ph/0303041].

[32] S. M. Carroll, V. Duvvuri, M. Trodden and M. S. Turner, "Is cosmic speed - up due to new gravitational physics?," Phys. Rev. D **70**, 043528 (2004) [arXiv:astro-ph/0306438].

[33] S. Nojiri and S. D. Odintsov, "Modified gravity with negative and positive powers of the curvature: Unification of the inflation and of the cosmic acceleration," Phys. Rev. D **68**, 123512 (2003) [arXiv:hep-th/0307288].